Supplementary Table 2. Age-adjusted incidence rate* of acute lymphoblastic leukemia,† aged <20 years, by state and race/ethnicity—United States, \$ 2001–2014

| State [¶] | White, non-Hispanic | | Black, non-Hispanic | | | Hispanic | |
|----------------------|---------------------|------------------|---------------------|------------------|-------|------------------|--|
| | No. | Rate (95% CI) | No. | Rate (95% CI) | No. | Rate (95% CI) | |
| Alaska | 48 | 28.4 (20.9–37.6) | _ | _ | _ | _ | |
| Alabama | 329 | 30.8 (27.6-34.3) | 98 | 18.0 (14.6-21.9) | 30 | 32.9 (22.0-47.3) | |
| Arkansas | _ | _ | _ | _ | _ | _ | |
| Arizona | 365 | 33.3 (30.0-36.9) | 18 | 15.0 (8.9-23.8) | 411 | 40.2 (36.4-44.2) | |
| California | 1,643 | 36.4 (34.7-38.3) | 217 | 22.0 (19.1-25.1) | 3,450 | 48.1 (46.5-49.7) | |
| Colorado | 400 | 34.8 (31.5-38.4) | _ | _ | 227 | 41.2 (36.0-46.9) | |
| Connecticut | 301 | 37.3 (33.2-41.7) | 16 | 10.1 (5.8-16.5) | 95 | 40.9 (33.1-50.0) | |
| District of Columbia | _ | _ | _ | _ | _ | _ | |
| Delaware | _ | _ | _ | _ | _ | _ | |
| Florida | 1,041 | 34.6 (32.5-36.7) | 236 | 17.8 (15.6-20.2) | 666 | 42.1 (39.0-45.4) | |
| Georgia | 641 | 33.9 (31.3-36.6) | 243 | 18.6 (16.3-21.1) | 202 | 45.4 (39.2-52.2) | |
| Hawaii | 27 | 44.5 (29.3-64.8) | _ | _ | _ | _ | |
| Idaho | 171 | 33.4 (28.6-38.8) | _ | _ | 39 | 38.5 (37.3-52.7) | |
| Illinois | 967 | 35.7 (33.5-38.1) | 151 | 17.5 (14.9-20.6) | 478 | 44.8 (40.9-49.0) | |
| Indiana | 664 | 34.9 (32.3-37.6) | 57 | 18.6 (14.1-24.1) | 99 | 48.0 (38.9-58.5) | |
| Iowa | 303 | 31.9 (28.4-35.7) | _ | _ | 35 | 38.6 (26.8-53.9) | |
| Kansas | _ | _ | _ | _ | 74 | 42.9 (33.6-54.0) | |
| Kentucky | _ | _ | _ | _ | _ | _ | |
| Louisiana | 305 | 31.8 (28.4-35.6) | 119 | 17.1 (14.2-20.5) | 36 | 43.4 (30.2-60.5) | |
| Massachusetts | _ | _ | _ | | _ | . , | |
| Maryland | 303 | 28.6 (25.5-32.0) | 111 | 15.6 (12.8-18.8) | 68 | 31.8 (24.6-40.4) | |
| Maine | 155 | 39.6 (33.6–46.3) | _ | _ | _ | _ | |
| Michigan | 913 | 34.7 (32.5–37.0) | 125 | 18.4 (15.3-21.9) | 79 | 31.0 (24.6-38.7) | |
| Minnesota | 519 | 33.8 (30.9–36.8) | 30 | 18.2 (12.2–26.0) | 61 | 41.9 (31.9–54.0) | |
| Missouri | 544 | 32.1 (29.5–34.9) | 76 | 22.4 (17.7–28.1) | 38 | 33.1 (23.3–45.6) | |
| Mississippi | _ | _ | _ | | _ | . , | |
| Montana | 79 | 27.8 (22.0-34.7) | _ | _ | _ | _ | |
| North Carolina | 653 | 32.7 (30.2–35.3) | 151 | 17.4 (14.7-20.4) | 193 | 46.5 (40.0-53.7) | |
| North Dakota | 70 | 34.1 (26.6–43.1) | _ | | _ | , , | |
| Nebraska | 160 | 29.5 (25.1–34.4) | _ | _ | 42 | 42.2 (30.2-57.3) | |
| New Hampshire | 162 | 39.8 (33.8–46.4) | _ | _ | _ | | |
| New Jersey | 673 | 38.8 (35.9–41.8) | 115 | 23.3 (19.2-27.9) | 247 | 35.8 (31.5-40.6) | |
| New Mexico | 101 | 45.0 (36.6–54.7) | _ | | 166 | 37.1 (31.7–43.2) | |
| Nevada | _ | _ | _ | _ | _ | , , | |
| New York | 1,362 | 37.5 (35.5-39.5) | 235 | 19.7 (17.2-22.4) | 570 | 37.8 (34.8-41.0) | |
| Ohio | 1,057 | 32.3 (30.4–34.3) | 108 | 15.5 (12.7–18.7) | 53 | 27.3 (20.4–35.9) | |
| Oklahoma | 290 | 32.7 (29.1–36.7) | 27 | 17.9 (11.8–26.1) | 79 | 42.5 (33.6–53.1) | |
| Oregon | 345 | 36.4 (32.7–40.5) | _ | | 121 | 47.0 (39.0–56.3) | |
| Pennsylvania | _ | _ | _ | _ | 130 | 35.3 (29.5–42.0) | |
| Rhode Island | 94 | 39.1 (31.5-47.8) | _ | _ | _ | , , | |
| South Carolina | 300 | 31.7 (28.2–35.4) | 89 | 15.8 (12.7-19.4) | 46 | 43.1 (31.3-57.7) | |
| South Dakota | 71 | 28.8 (22.5–36.3) | _ | . , | _ | , , | |
| Tennessee | 544 | 34.1 (31.3–37.1) | 109 | 22.2 (18.2-26.8) | 65 | 44.4 (34.0-57.0) | |
| Texas | 1,259 | 33.1 (31.3–35.0) | 253 | 19.8 (17.4–22.3) | 2,082 | 43.0 (41.1–44.9) | |
| Utah | 337 | 32.4 (29.0–36.0) | _ | _ | 96 | 48.5 (39.2–59.4) | |
| Virginia | 535 | 31.0 (28.4–33.7) | 118 | 18.1 (15.0-21.7) | 114 | 39.6 (32.6–47.7) | |
| Vermont | 85 | 44.3 (35.3–54.8) | _ | _ | _ | - | |
| Washington | 569 | 35.5 (32.6–38.5) | 39 | 27.9 (19.8-38.1) | 180 | 42.4 (36.4–49.2) | |
| Wisconsin | 533 | 34.0 (31.1–37.0) | 38 | 18.6 (13.2–25.5) | 74 | 39.7 (31.1–50.0) | |
| West Virginia | 164 | 29.5 (25.2–34.4) | _ | _ | _ | _ | |
| Wyoming | 42 | 25.1 (18.1–33.9) | _ | _ | _ | _ | |

Source: CDC's National Program of Cancer Registries and the National Cancer Institute's Surveillance, Epidemiology, and End Results program

Abbreviations: CI = confidence interval.

^{*} Rates are per 1 million persons and age-adjusted to the 2000 U.S. standard population. Rate statistics were not displayed if case count was <16 cases.

[†] Cases included International Classification of Diseases for Oncology, Third Edition codes (9728–9729, 9811–9818, 9835–9837) as grouped by the International Classification of Childhood Cancer.

⁹ Incidence data are compiled from cancer registries that meet the data quality criteria for all years 2001–2014 (covering approximately 98% of the U.S. population). Registry-specific data quality information is available at https://www.cdc.gov/cancer/npcr/uscs/data/00_data_quality.htm.

[¶] Race/ethnicity data was suppressed for states that elected to be excluded from race/ethnity analysis: Arkansas, Delaware, Kentucky, and Massachusetts. Kansas and Pennsylvania were excluded from presentation of rates by non-Hispanic ethnicities only.